OPEN ACCESS



The Journal of Threatened Taxa is dedicated to building evidence for conservation globally by publishing peer-reviewed articles online every month at a reasonably rapid rate at www.threatenedtaxa.org. All articles published in JoTT are registered under Creative Commons Attribution 4.0 International License unless otherwise mentioned. JoTT allows unrestricted use of articles in any medium, reproduction, and distribution by providing adequate credit to the authors and the source of publication.



Journal of Threatened Taxa

Building evidence for conservation globally

www.threatenedtaxa.org

ISSN 0974-7907 (Online) | ISSN 0974-7893 (Print)

Note

A RECORD OF *LIMENITIS RILEYI* TYTLER, 1940 (LEPIDOPTERA, NYMPHALIDAE, LIMENITIDINAE) FROM ARUNACHAL PRADESH, INDIA

Purnendu Roy

26 January 2017 | Vol. 9| No. 1 | Pp. 9774–9776 10.11609/jott.3070.9.1.9774-9776



For Focus, Scope, Aims, Policies and Guidelines visit http://threatenedtaxa.org/About_JoTT.asp
For Article Submission Guidelines visit http://threatenedtaxa.org/Submission_Guidelines.asp
For Policies against Scientific Misconduct visit http://threatenedtaxa.org/JoTT_Policy_against_Scientific_Misconduct.asp
For reprints contact <info@threatenedtaxa.org>

Partner



Publisher/Host





ISSN 0974-7907 (Online) ISSN 0974-7893 (Print)

OPEN ACCESS



Limenitis rileyi Tytler, 1940 (image 2) was described as a subspecies of Limenitis oberthuri Leech, 1890 from Hthawgaw (Fig. 1, site 4), northeastern Myanmar, June. Monastyrskii & Devyatkin (2000) raised L. rileyi to species status. Huang (2003) illustrated the genitalia for the two species and also published a checklist of the

subspecies. Monastyrskii & Devyatkin (2000) clarified that *Limenitis oberthuri* Leech, 1890 is a synonym of *Limenitis mimica* Poujade, 1885 (image 4). Monastyrskii & Devyatkin (2000) also clarified that the name *safeda* used in one of Tytler specimens (image 2) was invalid as it was an unpublished name. The species has previously been recorded from southeastern Tibet, Myanmar and northern Vietnam.

A single male specimen of *L. rileyi* (image1) was collected by the author on 19 July 1987 (approx. 1,800m) near Anini, Upper Dibang Valley District, Arunachal Pradesh, India (Fig. 1, site 3) in wet sub-tropical broad leaf forest.

A neighbouring subspecies *L. rileyi xizangana* Huang, 1998 (Image 3) has been recorded in southeastern Tibet (Yigong, Yigong Tsangpo (Fig. 1, site 1) and Metok, Lower Tsangpo (Fig. 1, site 2). The Dibang Valley record lies between the southeastern Tibet and northeastern Myanmar records and as such fills a gap in the distribution. No subspecies determination has been made, given the sample size of just one specimen.

There are no similar species in India, however both *L. mimica and L. rileyi* have been recorded near Kangfang (Fig. 1, site 5), Kachin state, Myanmar (Shizuya et al.

A RECORD OF *LIMENITIS RILEYI* TYTLER, 1940 (LEPIDOPTERA, NYMPHALIDAE, LIMENITIDINAE) FROM ARUNACHAL PRADESH, INDIA

Purnendu Roy

3 Gabriel's Wharf, 56 Upper Ground, London SE1 9PP purnendu@ganesha.co.uk

2011), so it is therefore possible that *L. mimica* may occur in India. The following morphological characters of *L. rileyi* will distinguish it from *L. mimica*:

- 1. Forewing cell completely filled with grey white in *L. rileyi*. In *L. mimica* the distal end of the cell has a grey-brown to grey-black triangle.
- 2. Under hindwing ground colour is reddishbrown in *L. rileyi*, not grey-brown as in *L. mimica*.
- 3. Upperside white markings generally wider between the veins in *L. rileyi*.

Records of *L. rileyi* from southeastern Tibet and northeastern Myanmar indicate a flight period of June to August at altitudes between 1,600–2,400 m. This sighting of *L. rileyi* represents the first record of this species from India, thus adding to India's butterfly fauna. This species is not listed in Varshney & Smetacek (2015). This species has the general appearance of a *Parantica/Neptis* species and as such could easily be overlooked, which may account for the lack of records of this species.

References

Huang (2003). A list of butterflies collected from Nujiang (Lou Tse Kiang) and Dulongiang, China with descriptions of new species, and revisional notes. *Neue Entomologische Nachrichten* 55: 3–114.

Monastyrskii & Devyatkin (2000). New taxa and new records of

DOI: http://doi.org/10.11609/jott.3070.9.1.9774-9776 | **ZooBank:** urn:lsid:zoobank.org:pub:EA326D1B-5079-4958-B94E-C5F2C5D00DA0

Editor: Anonymity requested. Date of publication: 26 January 2017 (online & print)

Manuscript details: Ms # 3070 | Received 27 September 2016 | Final received 01 January 2017 | Finally accepted 05 January 2017

Citation: Roy, P. (2017). A record of *Limenitis rileyi* Tytler, 1940 (Lepidoptera, Nymphalidae, Limenitidinae) from Arunachal Pradesh, India. *Journal of Threatened Taxa* 9(1): 9774–9776; http://doi.org/10.11609/jott.3070.9.1.9774-9776

Copyright: © Roy 2017. Creative Commons Attribution 4.0 International License. JoTT allows unrestricted use of this article in any medium, reproduction and distribution by providing adequate credit to the authors and the source of publication.

Funding: None.

 $\label{lem:competing} \textbf{Competing interests:} \ \ \textbf{The author declares no competing interests.}$

Acknowledgements: Tipa Umbrey and all the people who guided me in the Dibang Valley. David Lees of the Natural History Museum, London for his generous help and assistance. Jo Lawbuary my partner for her love and support.




Figure 1. Records of *L. riley*i from India and neighbouring countries.



Image 1. Limenitis rileyi (Tytler, 1940) male #982925 (left dorsal, right ventral) Dri Valley, Upper Dibang Valley, Arunachal Pradesh, c.1,830m, 19th July 1987



Image 2. Limenitis rileyi rileyi (Tytler, 1940) male paratype #982996 (left dorsal, right ventral) Htawgaw northeastern Myanmar, 1,500–2,400m, 25th June 1929







Image 3. Limenitis rileyi xizangana (Huang, 1998) male paratype #982999 (left dorsal, right ventral) Yigong southeastern Tibet, 2,400m, 4th Aug 1996







Image 4. Limenitis mimica (Poujade, 1885) (specimen shown is the male holotype of oberthuri Leech,1890) #982997 (left dorsal, right ventral) ChangYang, China, June 1888.

Photographs © Trustees of the Natural History Museum, London; used with permission (Original photography by David Lees)

butterflies from Vietnam. Atalanta 31(3/4): 471–492.

Shizuya, H., Y. Watanabe, M. Saito, T. Soe & H. Ozawa (2011) Basic information on butterflies of Kachin State, Myanmar (Part 5). Journal of Butterflies (Teinopalpus) 11–26.

Tytler, H.C. (1940). Notes on some new and interesting butterflies chiefly from Burma. Part 2. Journal of the Bombay Natural History Society 42: 109-123.

Varshney, R.K. & P. Smetacek (eds.) (2015). A Synoptic Catalogue of the Butterflies of India. Butterfly Research Centre, Bhimtal and Indinov Publishing, New Delhi, ii+261pp+8pls.







OPEN ACCESS The Journal of Threatened Taxa is dedicated to building evidence for conservation globally by publishing peer-reviewed articles online every month at a reasonably rapid rate at www.threatenedtaxa.org. All articles published in JoTT are registered under Creative Commons Attribution 4.0 International License unless otherwise mentioned. JoTT allows unrestricted use of articles in any medium, reproduction, and distribution by providing adequate credit to the authors and the source of publication.

ISSN 0974-7907 (Online); ISSN 0974-7893 (Print)

January 2017 | Vol. 9 | No. 1 | Pages: 9689-9776 Date of Publication: 26 January 2017 (Online & Print) DOI: 10.11609/jott.2017.9.1.9689-9776

www.threatenedtaxa.org

Article

Influence of substrate features on distribution of polypores (Fungi: Basidiomycota) in central part of Peechi Vazhani Wildlife Sanctuary, Kerala, India

-- Muhammed Iqbal, Kattany Vidyasagaran & Narayan Ganesh, Pp. 9689-9699

Reviews

Nepal's National Red List of Birds

-- Carol Inskipp, Hem Sagar Baral, Tim Inskipp, Ambika Prasad Khatiwada, Monsoon Pokharel Khatiwada, Laxman Prasad Poudyal & Rajan Amin, Pp. 9700-9722

Current status, distribution and conservation status of Algerian bats (Mammalia: Chiroptera)

-- Mourad Ahmim, Pp. 9723-9733

Short Communications

Incipient loss of a rainforest mutualism?

-- Johannes H. Fischer, Heiko U. Wittmer, Endro Setiawan, Sarah Jaffe & Andrew J. Marshall, Pp. 9734-9737

First record of the Woolly-necked Stork Ciconia episcopus Boddaert, 1783 (Aves: Ciconiiformes: Ciconiidae) breeding in the lowland wet zone of Sri Lanka

-- Hemachandra Kularatne & Susantha Udagedara, Pp. 9738-9742

Bibliography and checklist of the dragonflies and damselflies of Bhutan

-- T. Gyeltshen, T. Nidup, P. Dorji, T. Dorji & V.J. Kalkman, Pp. 9743-9747

Checklist of terebrantian thrips (Insecta: Thysanoptera) recorded from India

-- R.R. Rachana & R. Varatharajan, Pp. 9748-9755

Notes

Two jasmine (Oleaceae: Jasminum L.) taxa newly recorded in Vietnam

-- Bui Hong Quang, Vu Tien Chinh, Le Thi Mai Linh & Ritesh Kumar Choudhary, Pp. 9756-9760

First record of Rusty-spotted Cat Prionailurus rubiginosus (Mammalia: Carnivora: Felidae) from Ramgarh-Vishdhari Wildlife Sanctuary in semi-arid landscape of Rajasthan, India -- Sailaja Nayak, Sunny Shah & Jimmy Borah, Pp. 9761-9763

Re-sighting record of Fulvous Leaf-nosed Bat Hipposideros fulvus Gray, 1838 (Mammalia: Chiroptera: Hipposideridae) from Thar Desert, Rajasthan, India

-- Sumit Dookia, Gajendra Singh & Rajlakshmi Mishra, Pp. 9764-9767

Record number of Yellow-billed Oxpeckers Buphagus africanus Linnaeus, 1766 (Aves: Passeriformes: Buphagidae) foraging on a single host

-- Diogo Veríssimo, Jean-Christophe Cugnière, Suzanne Cugnière, Julien Cugnière, Géraldine Cugnière & Laure Cugnière, Pp. 9768-9770

First record of the Two-striped Box Crab Calappa bilineata Ng, Lai & Aungtonya, 2002 (Brachyura: Calappidae) from St. Martin's Island, Bangladesh

-- Muntasir Akash & Mostafa A.R. Hossain, Pp. 9771–9773

A record of Limenitis rileyi Tytler, 1940 (Lepidoptera, Nymphalidae, Limenitidinae) from Arunachal Pradesh, India

-- Purnendu Roy, Pp. 9774-9776



